

1003 526

CLENSON UNIV S C DEPT OF MATHEMATICAL SCIENCES

F/8 12/1

THE INITIAL BUSY CYCLE OF A DISCRETE-TIME G1/G/1 QUEUE IN WHICH-ETC(U)

JUN 79 D L MINH


N00014-75-C-0451

UNCLASSIFIED

N110

NL

1 of 1  
AD A  
083026




END  
DATE  
FILMED  
5-80  
DTIC

12


LEVEL II

00  
31  
10  
03  
80  
11



DEPARTMENT  
OF  
MATHEMATICAL  
SCIENCES

CLEMSON UNIVERSITY  
Clemson, South Carolina



FILE COPY

DTIC

S

DTIC  
ELECTE  
APR 28 1980

D

B

DISTRIBUTION STATEMENT A

Approved for public release;  
Distribution Unlimited

80 4 24 0 11

12

LEVEL

THE INITIAL BUSY CYCLE OF A DISCRETE-  
TIME GI/G/1 QUEUE IN WHICH THE  
INTER-ARRIVAL TIMES ARE NOT  
NECESSARILY IDENTICALLY DISTRIBUTED.

Do Le Minh\*

Clemson University

Technical Report #319

June 1979

\*On leave from The New South Wales Institute of Technology, P.O. Box 123,  
Broadway, NSW 2007, Australia.

Research supported in part by THE OFFICE OF NAVAL RESEARCH Contract  
N00014-75-C-0451

DISTRIBUTION STATEMENT A  
Approved for public release  
Distribution Unlimited

DTIC  
SELECTED  
APR 20 1980

B

THE INITIAL BUSY CYCLE OF A DISCRETE-TIME GI/G/1 QUEUE IN WHICH  
THE INTER-ARRIVAL TIMES ARE NOT NECESSARILY IDENTICALLY DISTRIBUTED.

Do Le Minh,  
Department of Mathematical Sciences,  
Clemson University,  
Clemson, SC 29631, U.S.A. (\*)

ABSTRACT.

This paper studies a generalization of the discrete-time GI/G/1 queueing system. Here, the inter-arrival times are not necessarily identically distributed. A recursive scheme is derived to obtain the joint distribution of the duration of the initial busy period, the duration of the first idle period and the number of customers served during the initial busy period.

KEY WORDS AND PHRASES: GI/G/1 queue. Inter-arrival times not identically distributed. Initial busy period. First idle period. Number of customers served during the initial busy period.

(\*) On Leave from The New South Wales Institute of Technology,  
P.O. Box 123, Broadway, NSW 2007, Australia.

(\*) Research supported in part by THE OFFICE OF NAVAL RESEARCH  
Contract N00014-75-C-0451.

Section	<input checked="checked" type="checkbox"/>
ction	<input type="checkbox"/>
	<input type="checkbox"/>
Y CODES	
SPECIAL	

A

The initial busy cycle of a queueing system has always been a subject of interest. It is of great importance in the study of systems in which the cost associated with unproductive idle period is high and thus the server has to be 'switch off' or transferred to another system whenever the original system is empty for the first time. Most of the results obtained in the literature for the initial busy cycle, however, are not very suitable for computer applications. For the GI/G/1 queue, they are presented in terms of Laplace-Stieljes transforms involving complicated operators ([8], equ. 5.18). For the M/G/1 queue, they contain the roots of a certain functional equation within an unit circle ([3], equ. 25). Obtaining numerical information from these results is not an easy task and has never been discussed thoroughly in the literature.

In this paper, we study a discrete-time single-server queueing system in which the inter-arrival times are mutually independent but not necessarily identically distributed; the service times are mutually independent and identically distributed. We shall develop a recursive scheme which enables us to calculate easily the joint distribution of the duration of the initial busy period, the duration of the first idle period and the number of customers served during the initial busy period. This scheme is distribution free; that is, it requires no specific distributional assumptions for the random variables underlying the model. This scheme is suitable

for systems having a very general initial condition. When the system is initiated by the arrival of a customer to an empty queue, this scheme will enable us to obtain information about the normal busy cycle. For the queueing system in which there is a random 'set-up' time for customers who arrive when the server is idle ([7], [3]) this scheme is also applicable as the set-up times only affect the initiation of the busy cycle.

For the actual waiting time of each customer in this model, another recursive scheme has been reported in [9].

Let us restrict ourselves to the discrete-time system observed at equally spaced epochs  $0, 1, 2, \dots, n, \dots$ . All events of the system namely arrivals, transfers from queue to service and departures are assumed to occur at instants immediately prior to these epochs. We assume that initially the system is not empty and that there are  $k_0$  ( $k_0 \geq 1$ ) customers in it at the epoch  $n=0$  and the  $(k_0+1)$ th,  $(k_0+2)$ th, ...,  $k$ th, ... customers arrive at the epochs  $\tau_{k_0+1}, \tau_{k_0+2}, \dots, \tau_k, \dots$  where  $0 < \tau_{k_0+1} < \tau_{k_0+2} < \dots < \tau_k < \dots$ . We also assume that the inter-arrival times  $t_k = \tau_{k+1} - \tau_k$  ( $k \geq k_0$ ) are mutually independent but not necessarily identically distributed. We write

$$(1) \quad a_v^k = \Pr(t_k = v), \quad v \geq 1, k \geq k_0; \quad \alpha_k(z) = \sum_{v=1}^{\infty} a_v^k z^v, \quad z \leq 1, k \geq k_0.$$

Let the service time of each customer be  $S$ . We write

$$(2) \quad s_j = \Pr(S=j), \quad j \geq 1; \quad \beta(z) = \sum_{j=1}^{\infty} s_j z^j, \quad z \leq 1.$$

The customers are served in order of their arrivals and there is no limit on the waiting room.

First, let  $\tilde{V}_n$  ( $n \geq 0$ ) be the virtual waiting time at epoch  $n$ . Let  $\tilde{A}_n$  ( $n \geq 0$ ) be the time difference between  $n$  and the time of the first arrival in the interval  $(n, \infty)$ . Let  $\tilde{C}_n$  ( $n \geq 0$ ) be the total number of customers being in the system at the epoch  $n=0$  or arriving at the system during the interval  $(0, n]$ . We shall study the multi-variate Markov chain  $\{\tilde{V}_n, \tilde{A}_n; \tilde{C}_n\}$ .

It is well-known that any stochastic process can be characterized as a Markov process if the full set of random variables needed to specify the state of the process is employed. For the queueing system studied in this paper, the instantaneous state of the process is completely characterized by the bi-variate  $\{\tilde{V}_n, \tilde{A}_n\}$ . In this paper, however, instead of studying the bi-variate  $\{\tilde{V}_n, \tilde{A}_n\}$ , we study the tri-variate  $\{\tilde{V}_n, \tilde{A}_n; \tilde{C}_n\}$ . We shall show that this will enable us to resolve our problem more fully (See also [6], [7], [9]).

Now let

$$(3) \quad v_{j, \ell; k}^n = \Pr\{\tilde{V}_n = j, \tilde{A}_n = \ell; \tilde{C}_n = k\} \quad (n \geq 0, j \geq 1, \ell \geq 1, k \geq 1).$$

Straightforward arguments result in the following Chapman-Kolmogorov difference equations for  $n \geq 0$ :

$$(4) \quad v_{0, \ell; k}^{n+1} = v_{0, \ell+1; k}^n + v_{1, \ell+1, k}^n \quad (\ell \geq 1, k \geq 1);$$

$$(5) \quad v_{j, \ell; 1}^{n+1} = v_{j+1, \ell+1; 1}^n \quad (j \geq 1, \ell \geq 1);$$

$$(6) \quad v_{j, \ell; k}^{n+1} = v_{j+1, \ell+1; k}^n + a_{\ell-m=1}^{k-j} v_{j-m+1, 1, k-1}^n s_m + v_{0, 1; k-1}^n s_j a_{\ell}^k \quad (j \geq 1, \ell \geq 1, k \geq 2).$$

Let us now consider a second queueing system, having exactly the same description as the one above but admitting no more customers whenever the system is empty for the first time. It is equivalent to the initial busy period of the original queueing system. The second system will have the same Chapman-Kolmogorov difference equations as (4)-(6) except that the term  $v_{0,1;k-1}^n s_j a_l^k$  does not appear in (6).

Let the notation for the second queueing system be those for the first system together with the additional symbol  $\hat{\cdot}$ . We say that the chain  $\{\hat{V}_n, \hat{A}_n, \hat{C}_n\}$  has  $\{0, 1; k \geq 1, k \geq 1\}$  as absorbing states. Cox and Miller [2, p.236] suggest this approach as one suitable method to investigate the distribution of the busy period of the M/G/1 model but do not elaborate any further. (See also [4] and [5]).

From (1), (2), (5)-(6), with the term  $v_{0,1;k-1}^n s_j a_l^k$  omitted, after some routine calculations, we obtain the following relation between the generating functions

$$\begin{aligned}
 (7) \quad & \sum_{j=1}^{\infty} \sum_{l=1}^{\infty} \hat{v}_{j,l;k}^{n+1} z^j t^l \\
 &= \frac{1}{zt} \sum_{j=1}^{\infty} \sum_{l=1}^{\infty} \hat{v}_{j,l;k}^n z^j t^l - \frac{1}{z} \sum_{j=1}^{\infty} \hat{v}_{j,1;k}^n z^j \\
 &\quad - \frac{1}{t} \sum_{l=2}^{\infty} \hat{v}_{1,l;k}^n t^l - (1 - \delta_{k1}) \delta(z) \delta_k(t) \sum_{j=1}^{\infty} \hat{v}_{j,1;k-1}^n z^j \\
 &\quad (0 < |z| \leq 1, 0 < |t| \leq 1, n \geq 0, k \geq 1)
 \end{aligned}$$

where  $\delta_{ij}$  is the Kronecker delta.

$\hat{v}_{1,l;k}^n$  ( $n \geq 0, l \geq 2, k \geq 1$ ) is the probability that the second queueing system becomes empty for the first time after the departure of the  $k^{\text{th}}$  customer at epoch  $n+1$  and that the  $(k-1)^{\text{th}}$  customer would have arrived at the system at epoch  $(n-1)$  were he allowed to



do so. It is therefore equal to the joint probability that the initial busy period of the original system ends at epoch  $(n+1)$ , there are  $k$  customers served during this period and the duration of the first idle period is  $(\ell-1)$ . If we denote this by  $b_{\ell-1;k}^{n+1}$ , we shall have

$$(8) \quad \xi \sum_{n=0}^{\infty} \sum_{\ell=2}^{\infty} \hat{v}_{1,\ell;k}^n t^{\ell} \xi^n = t \sum_{n=1}^{\infty} \sum_{\ell=1}^{\infty} b_{\ell;k}^n t^{\ell} \xi^n \quad (|t| \leq 1, |\xi| \leq 1, k \geq 1).$$

This relationship indicates that the busy period studied in this paper is what Cohen [1, p. 284] calls the *strong* busy period. A strong busy period continues when an arrival occurs at the instant of departure of the last customer in the system. In other words, two strong busy periods are separated with probability one by an idle period of non-zero duration.

From (7), taking the generating function with respect to time yields

$$\begin{aligned} (9) \quad & \left[ \frac{1}{\xi} - \frac{1}{zt} \right] \sum_{n=0}^{\infty} \sum_{j=1}^{\infty} \sum_{\ell=1}^{\infty} \hat{v}_{j,\ell;k}^n z^j t^{\ell} \xi^n \\ &= \frac{1}{\xi} \sum_{j=1}^{\infty} \sum_{\ell=1}^{\infty} \hat{v}_{j,\ell;k}^0 z^j t^{\ell} - \frac{1}{z} \sum_{n=0}^{\infty} \sum_{j=1}^{\infty} \hat{v}_{j,1;k}^n z^j \xi^n \\ &\quad - \frac{1}{\xi} \sum_{n=1}^{\infty} \sum_{\ell=1}^{\infty} b_{\ell;k}^n t^{\ell} \xi^n \\ &\quad + (1-\delta_{k1}) \Psi(z) \beta_k(t) \frac{1}{z} \sum_{n=0}^{\infty} \sum_{j=1}^{\infty} \hat{v}_{j,1,k-1}^n z^j \xi^n \\ &\quad (0 < |z| \leq 1, 0 < |t| \leq 1, |\xi| \leq 1, k \geq 1). \end{aligned}$$

If  $0 < |\xi| \leq |t| \leq 1$ ,  $k \geq 1$ , letting  $z = \xi/t$  in the above equation we obtain

$$(10) \quad \sum_{n=0}^{\infty} \sum_{j=1}^{\infty} \hat{v}_{j,1;k}^n (\xi/t)^j \xi^n + \frac{1}{t} \sum_{n=1}^{\infty} \sum_{l=1}^{\infty} b_{l;k}^n t^l \xi^n$$

$$= \frac{1}{t} \sum_{j=1}^{\infty} \sum_{l=1}^{\infty} \hat{v}_{j,l;k}^0 (\xi/t)^j t^l$$

$$+ (1 - \delta_{k1}) \Psi(\xi/t) \Omega_k(t) \sum_{n=0}^{\infty} \sum_{j=1}^{\infty} \hat{v}_{j,1;k-1}^n (\xi/t)^j \xi^n.$$

From (10), denoting by  $\beta_u^{(v)}$  the coefficient of  $z^u$  in  $\psi^v(z)$  and by  $a_u^{m,n}$  the coefficient of  $z^u$  in  $\Omega_{v=m}^n(z)$ , by induction, we get

$$(11) \quad t \sum_{n=0}^{\infty} \sum_{j=1}^{\infty} \hat{v}_{j,1;k}^n t^{-j} \xi^{j+n} + \sum_{n=1}^{\infty} \sum_{l=1}^{\infty} b_{l;k}^n t^l \xi^n$$

$$= \sum_{j=1}^{\infty} \sum_{l=1}^{\infty} \hat{v}_{j,l;k}^0 t^{l-j} \xi^j$$

$$+ \sum_{v=1}^{k-1} \left[ \sum_{u=1}^{\infty} \beta_u^{(k-v)} \xi^u t^{-u} \right] \left[ \sum_{u=1}^{\infty} a_u^{v+1,k} t^u \right] \sum_{j=1}^{\infty} \sum_{l=1}^{\infty} \hat{v}_{j,l;v}^0 t^{l-j} \xi^j$$

$$- \sum_{v=1}^{k-1} \left[ \sum_{u=1}^{\infty} \beta_u^{(k-v)} \xi^u t^{-u} \right] \left[ \sum_{u=1}^{\infty} a_u^{v+1,k} t^u \right] \sum_{n=1}^{\infty} \sum_{l=1}^{\infty} b_{l;v}^n t^l \xi^n$$

$$(0 < |\xi| \leq |t| \leq 1, k \geq 2).$$

Equating the positive-degree coefficients of  $t$  in the above equation yields

$$(12) \quad \sum_{n=1}^{\infty} b_{l;k}^n \xi^n = \sum_{j=1}^{\infty} \hat{v}_{j,l+j;k}^0 \xi^j$$

$$+ \sum_{v=1}^{k-1} \sum_{u=1}^{\infty} \sum_{w=-j}^{u+l} \beta_{u-w}^{(k-v)} a_{u-w+1,k-v}^{v+1,k} \hat{v}_{j,j+w;v}^0 \xi^{j+u}$$

$$= \sum_{v=1}^{k-1} \sum_{u=1}^{\infty} \sum_{w=1}^{u+\lambda} 3_u^{(k-v)} a_{u-w+\lambda}^{v+1, k} \sum_{m=1}^{\infty} b_{w, v}^m \xi^{m+u}$$

$$(0 < |\xi| < 1, \lambda \geq 1, k \geq 2).$$

Further, equating the coefficients of  $\xi$  yields the following relations which will allow us to calculate easily the values of  $b_{\lambda; k}^n$  ( $n \geq 1, \lambda \geq 1, k \geq 2$ ) from the given initial distribution  $\hat{v}_{j, \lambda; k}^0 = v_{j, \lambda; k}^0 = \Pr\{V_0=j, A_0=\lambda; C_0=k_0=k\}$ :

$$\begin{aligned} (13) \quad b_{\lambda; k}^n &= v_{n, \lambda+n; k}^0 \\ &+ \sum_{v=1}^{k-1} \sum_{u=1}^n \sum_{w=u-n}^{u+\lambda} 3_u^{(k-v)} a_{u-w+\lambda}^{v+1, k} v_{n-u, n-u+w; v}^0 \\ &- \sum_{v=1}^{k-1} \sum_{u=1}^n \sum_{w=1}^{u+\lambda} 3_u^{(k-v)} a_{u-w+\lambda}^{v+1, k} b_{w, v}^{n-u} \end{aligned}$$

$$(n \geq 1, \lambda \geq 1, k \geq 2).$$

The recurrence is initiated by  $b_{\lambda; 1}^n = v_{n, \lambda+n; 1}^0$  ( $n \geq 1, \lambda \geq 1$ ).

If the initial busy period is initiated by the arrival of one customer to an empty queue, then  $v_{1, \lambda; k}^0 = \lambda_1 a_{\lambda}^1 s_1$  and (13) would enable us to calculate the joint distribution of the duration of a normal busy period, the duration of a normal idle period and the number of customers served during a normal busy period.

For the system in which there is a random set-up time  $H$  ( $H > 0$ ) for customers who arrive when the server is idle, we write  $h_1 = \Pr\{H=1 | H > 0\}$  and  $v_{1, \lambda; k}^0 = (\lambda_1 a_{\lambda}^{1-1} s_1)_{\lambda=1} h_1$  to obtain information about its normal busy cycle.

# REFERENCES:

1. J.W. Cohen, The single-server queue, North-Holland, Amsterdam (1969).
2. D.R. Cox and H.D. Miller, The Theory of Stochastic Processes, Methuen, London (1965).
3. S. Erlander, The remaining busy period for a single-server queue with Poisson input, Operations Res., 13, 734-746 (1965)
4. P. Hokstad, A supplementary variable technique applied to the M/G/1 queue, Scand. J. Statist., 2, 95-98 (1975)
5. P. Hokstad, The use of Wiener-Hopf decomposition in the study of waiting time and busy period distribution for the GI/G/1 queue, Scand. J. Statist., 3, 79-85 (1976).
6. D.L. Minh, The discrete-time, single-server queue with time-inhomogeneous compound Poisson input and general service time distribution, J. Appl. Prob., 15, 590-601 (1978).
7. D.L. Minh, A discrete-time single-server queue with set-up time, Stochastic Processes Appli., 3, 131-197 (1978).
8. D.L. Minh, Analysis of the exceptional queueing system by the use of regenerative processes and analytical methods, Maths Operat. Res., 24(3) (1979)
9. D.L. Minh, The actual waiting time of each customer in a GI/G/1 queue, J. Appl. Prob., 16, (1979)

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER N110	2. GOVT ACCESSION NO. AD AC 83 524	3. RECIPIENT'S CATALOG NUMBER	
4. TITLE and Subtitle The Initial Busy Cycle of a Discrete-Time GI/G/1 Queue in Which the Inter-Arrival Times Are Not Necessarily Identically Distributed.		5. TYPE OF REPORT & PERIOD COVERED	
7. AUTHOR(s) Do Le Minh		6. PERFORMING ORG. REPORT NUMBER Tech. Report #319	
		8. CONTRACT OR GRANT NUMBER(s) N00014-75-C-0451	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Clemson University Dept. of Mathematical Sciences Clemson, South Carolina 29631		10. PROGRAM ELEMENT PROJECT TASK AREA & WORK UNIT NUMBERS NR 042-271	
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Code 436 Arlington, Va. 22217		12. REPORT DATE June, 1979	
		13. NUMBER OF PAGES 9	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. of this report Unclassified	
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.			
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)			
18. SUPPLEMENTARY NOTES			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  GI/G/1 queue. Inter-arrival times not identically distributed. Initial busy period. First idle period. Number of customers served during the initial busy period.			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  This paper studies a generalization of the discrete-time GI/G/1 queueing system. Here, the inter-arrival times are not necessarily identically distributed. A recursive scheme is derived to obtain the joint distribution of the duration of the initial busy period, the duration of the first idle period and the number of customers served during the initial busy period.			

DD FORM 1473

EDITION OF 1 NOV 65 IS OBSOLETE  
S/N 1112-114-5601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

DATE  
FILMED  
— 8